

Datasheet for ABIN1881429

anti-HOXC10 antibody (C-Term)

2 Images

5 Publications

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	HOXC10
Binding Specificity:	AA 224-250, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HOXC10 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details

Immunogen:	This HOXC10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 224-250 amino acids from the C-terminal region of human HOXC10.
Clone:	RB42429
Isotype:	Ig Fraction
Predicted Reactivity:	M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	HOXC10
Alternative Name:	HOXC10 (HOXC10 Products)

Target Details

Background: This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conserved family of transcription factors that play an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, which are located on different chromosomes and consist of 9 to 11 genes arranged in tandem. This gene is one of several homeobox HOXC genes located in a cluster on chromosome 12. The protein level is controlled during cell differentiation and proliferation, which may indicate this protein has a role in origin activation.

Molecular Weight: 38073

NCBI Accession: [NP_059105](#)

UniProt: [Q9NYD6](#)

Application Details

Application Notes: IF: 1:10~50. WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications


Product cited in: Carrascal, Ovelleiro, Casas, Gay, Abian: "Phosphorylation analysis of primary human T lymphocytes using sequential IMAC and titanium oxide enrichment." in: **Journal of proteome research**, Vol. 7, Issue 12, pp. 5167-76, (2009) ([PubMed](#)).

Koulich, Li, DeMartino: "Relative structural and functional roles of multiple deubiquitylating proteins associated with mammalian 26S proteasome." in: **Molecular biology of the cell**, Vol.

19, Issue 3, pp. 1072-82, (2008) ([PubMed](#)).


Reuter, Medhurst, Waisfisz, Zhi, Herterich, Hoehn, Gross, Joenje, Hoatlin, Mathew, Huber: "Yeast two-hybrid screens imply involvement of Fanconi anemia proteins in transcription regulation, cell signaling, oxidative metabolism, and cellular transport." in: **Experimental cell research**, Vol. 289, Issue 2, pp. 211-21, (2003) ([PubMed](#)).

Images



Immunofluorescence

Image 1. Fluorescent image of HeLa cell stained with HOXC10 Antibody (C-term) (ABIN1881429 and ABIN2838971). HeLa cells were fixed with 4 % PFA (20 min), permeabilized with Triton X-100 (0.1 %, 10 min), then incubated with HOXC10 primary antibody (1:25, 1 h at 37 °C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37 °C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/mL, 1 h at 37 °C). HOXC10 immunoreactivity is localized to Nucleus significantly.



Western Blotting

Image 2. HOXC10 Antibody (C-term) (ABIN1881429 and ABIN2838971) western blot analysis in MDA-M cell line lysates (35 µg/lane). This demonstrates the HOXC10 antibody detected the HOXC10 protein (arrow).