

Datasheet for ABIN1881459
anti-ILKAP antibody (AA 93-122)[Go to Product page](#)

1 Image

3 Publications

Overview

Quantity:	400 µL
Target:	ILKAP
Binding Specificity:	AA 93-122
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ILKAP antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This ILKAP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 93-122 amino acids from the Central region of human ILKAP.
Clone:	RB40435
Isotype:	Ig Fraction
Predicted Reactivity:	B, M, Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	ILKAP
Alternative Name:	ILKAP (ILKAP Products)

Target Details

Background: The protein encoded by this gene is a protein serine/threonine phosphatase of the PP2C family. This protein can interact with integrin-linked kinase (ILK/ILK1), a regulator of integrin mediated signaling, and regulate the kinase activity of ILK. Through the interaction with ILK, this protein may selectively affect the signaling process of ILK-mediated glycogen synthase kinase 3 beta (GSK3beta), and thus participate in Wnt signaling pathway.

Molecular Weight: 42907

NCBI Accession: [NP_110395](#)

UniProt: [Q9H0C8](#)

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Expiry Date: 6 months

Publications

Product cited in: Hyrskyluoto, Bruelle, Lundh, Do, Kivinen, Rappou, Reijonen, Waltimo, Petersén, Lindholm, Korhonen: "Ubiquitin-specific protease-14 reduces cellular aggregates and protects against mutant huntingtin-induced cell degeneration: involvement of the proteasome and ER stress-activated kinase IRE1?." in: **Human molecular genetics**, Vol. 23, Issue 22, pp. 5928-39, (2014) ([PubMed](#)).

Davila, Froeling, Tan, Bonnard, Boland, Snippe, Hibberd, Seielstad: "New genetic associations detected in a host response study to hepatitis B vaccine." in: **Genes and immunity**, Vol. 11,


Issue 3, pp. 232-8, (2010) ([PubMed](#)).

Chen, Qin, Li, Walters, Wilson, Mei, Wilson: "The proteasome-associated deubiquitinating enzyme Usp14 is essential for the maintenance of synaptic ubiquitin levels and the development of neuromuscular junctions." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 29, Issue 35, pp. 10909-19, (2009) ([PubMed](#)).

Nagai, Kadowaki, Maruyama, Takeda, Nishitoh, Ichijo: "USP14 inhibits ER-associated degradation via interaction with IRE1alpha." in: **Biochemical and biophysical research communications**, Vol. 379, Issue 4, pp. 995-1000, (2009) ([PubMed](#)).

Mines, Goodwin, Limbird, Cui, Fan: "Deubiquitination of CXCR4 by USP14 is critical for both CXCL12-induced CXCR4 degradation and chemotaxis but not ERK activation." in: **The Journal of biological chemistry**, Vol. 284, Issue 9, pp. 5742-52, (2009) ([PubMed](#)).

Images



Western Blotting

Image 1. ILK Antibody (Center) 19399c western blot analysis in cell line lysates (35 µg/lane). This demonstrates the ILK antibody detected the ILK protein (arrow).