

Datasheet for ABIN1881491
anti-LATS2 antibody (C-Term)[Go to Product page](#)

2 Images

4 Publications

Overview

Quantity:	400 µL
Target:	LATS2
Binding Specificity:	AA 1000-1027, C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LATS2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This Mouse Lats2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1000-1027 amino acids from the C-terminal region of mouse Lats2.
Clone:	RB40566
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	LATS2
Alternative Name:	Lats2 (LATS2 Products)
Background:	Negative regulator of YAP1 in the Hippo signaling pathway that plays a pivotal role in organ size

Target Details

control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Phosphorylation of YAP1 by LATS2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. Acts as a tumor suppressor which plays a critical role in centrosome duplication, maintenance of mitotic fidelity and genomic stability. Negatively regulates G1/S transition by down-regulating cyclin E/CDK2 kinase activity. Negative regulator of the androgen receptor (By similarity).

Molecular Weight: 115472

NCBI Accession: [NP_056586](#)

UniProt: [Q7TSJ6](#)

Application Details

Application Notes: WB: 1:2000. IHC-P: 1:100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications

Product cited in: Dai, Liu, Liu, Zhang, Wang, Jin, Qian, Wang, Zhao, Wu, Xiong, Chang, Sun, Yang, Hoffman, Liu: "Anti-metastatic Efficacy of Traditional Chinese Medicine (TCM) Ginsenoside Conjugated to a VEGFR-3 Antibody on Human Gastric Cancer in an Orthotopic Mouse Model." in: **Anticancer research**, Vol. 37, Issue 3, pp. 979-986, (2017) ([PubMed](#)).

Irrthum, Karkkainen, Devriendt, Alitalo, Vikkula: "Congenital hereditary lymphedema caused by a mutation that inactivates VEGFR3 tyrosine kinase." in: **American journal of human genetics**, Vol. 67, Issue 2, pp. 295-301, (2000) ([PubMed](#)).

Galland, Karamysheva, Pebusque, Borg, Rottapel, Dubreuil, Rosnet, Birnbaum: "The FLT4 gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor." in: **Oncogene**, Vol. 8, Issue 5, pp. 1233-40, (1993) ([PubMed](#)).

Pajusola, Aprelikova, Korhonen, Kaipainen, Pertovaara, Alitalo, Alitalo: "FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines." in: **Cancer research**, Vol. 52, Issue 20, pp. 5738-43, (1992) ([PubMed](#)).


Galland, Karamysheva, Mattei, Rosnet, Marchetto, Birnbaum: "Chromosomal localization of FLT4, a novel receptor-type tyrosine kinase gene." in: **Genomics**, Vol. 13, Issue 2, pp. 475-8, (1992) ([PubMed](#)).

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of B on paraffin-embedded Mouse testis tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH 9.0). Samples were incubated with primary Antibody (1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Western Blotting

Image 2. All lanes : Anti-Mouse Lats2 Antibody (C-term) at 1:2000 dilution Lane 1: Mouse ovary tissue lysate Lane 2: Mouse testis tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 115 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.