



[Go to Product page](#)

Datasheet for ABIN1881495
anti-LGMN antibody (N-Term)

2 Images

3 Publications

Overview

Quantity:	400 µL
Target:	LGMN
Binding Specificity:	AA 83-112, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LGMN antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This LGMN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 83-112 amino acids from the N-terminal region of human LGMN.
Clone:	RB41868
Isotype:	Ig Fraction
Predicted Reactivity:	Pr
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	LGMN
Alternative Name:	LGMN (LGMN Products)

Target Details

Background: This gene encodes a cysteine protease that has a strict specificity for hydrolysis of asparaginyl bonds. This enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and appears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A fully mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. Overexpression of this gene may be associated with the majority of solid tumor types. This gene has a pseudogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two variants encode the same isoform.

Molecular Weight: 49411

NCBI Accession: [NP_001008530](#), [NP_005597](#)

UniProt: [Q99538](#)

Pathways: [Metabolism of Steroid Hormones and Vitamin D](#), [Activation of Innate immune Response](#), [Toll-Like Receptors Cascades](#)

Application Details

Application Notes: WB: 1:1000. WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications

Product cited in: Mehta, Vazquez, Kulkarni, Kerrigan, Atwal, Metsugi, Toppmeyer, Levine, Hirshfield: "Polymorphic

variants in TSC1 and TSC2 and their association with breast cancer phenotypes." in: **Breast cancer research and treatment**, Vol. 125, Issue 3, pp. 861-8, (2011) ([PubMed](#)).


Hoogeveen-Westerveld, Exalto, Maat-Kievit, van den Ouweland, Halley, Nellist: "Analysis of TSC1 truncations defines regions involved in TSC1 stability, aggregation and interaction." in: **Biochimica et biophysica acta**, Vol. 1802, Issue 9, pp. 774-81, (2010) ([PubMed](#)).

Mieulet, Lamb: "Tuberous sclerosis complex: linking cancer to metabolism." in: **Trends in molecular medicine**, Vol. 16, Issue 7, pp. 329-35, (2010) ([PubMed](#)).

Guo, Ying, Zhang, Yuan, Qian, Wang, Yang, He: "Tandem affinity purification and identification of the human TSC1 protein complex." in: **Acta biochimica et biophysica Sinica**, Vol. 42, Issue 4, pp. 266-73, (2010) ([PubMed](#)).


Liu, Wu, Chen, Ter-Minassian, Asomaning, Zhai, Wang, Su, Heist, Kulke, Lin, Liu, Christiani: "A Large-scale genetic association study of esophageal adenocarcinoma risk." in: **Carcinogenesis**, Vol. 31, Issue 7, pp. 1259-63, (2010) ([PubMed](#)).

Images



Western Blotting

Image 1. Anti-LGMN Antibody (N-term) at 1:1000 dilution + human kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa
Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 2. LGMN Antibody (N-term) (ABIN1881495 and ABIN2838891) western blot analysis in 293 cell line lysates (35 µg/lane). This demonstrates the LGMN antibody detected the LGMN protein (arrow).