antibodies .- online.com





anti-LIMK2 antibody (C-Term)

2 Images



Go to Product page

Overview	
Quantity:	400 μL
Target:	LIMK2
Binding Specificity:	AA 570-599, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LIMK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This LIMK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 570-599 amino acids from the C-terminal region of human LIMK2.
Clone:	RB42813
Isotype:	lg Fraction
Predicted Reactivity:	В
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	LIMK2
Alternative Name:	LIMK2 (LIMK2 Products)

Target Details

Background:	Displays serine/threonine-specific phosphorylation of myelin basic protein and histone (MBP) in vitro.
Molecular Weight:	72232
NCBI Accession:	NP_001026971, NP_005560, NP_057952
UniProt:	P53671


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:25
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffinembedded H.kidney section using LIMK2 Antibody (C-term) (ABIN1881499 and ABIN2843267). (ABIN1881499 and ABIN2843267) was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Western Blotting

Image 2. Western blot analysis of lysates from Jurkat cell line and human placenta tissue lysate (from left to right), using LIMK2 Antibody (C-term) (ABIN1881499 and ABIN2843267). (ABIN1881499 and ABIN2843267) was diluted at 1:1000 at each lane. A goat anti-rabbit $lgGH_{L}(HRP)$ at 1:5000 dilution was used as the secondary antibody. Lysate at 35 μ g per lane.