antibodies - online.com







anti-LRMP antibody (C-Term)





\sim	
()\/△	rview
\cup	1 410 44

Quantity:	400 μL
Target:	LRMP
Binding Specificity:	AA 403-431, C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRMP antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This Mouse Lrmp antibody is generated from rabbits immunized with a KLH conjugated
Immunogen:	This Mouse Lrmp antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp.
Immunogen: Clone:	
	synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp.
Clone:	synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp. RB40716
Clone:	synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp. RB40716 IgG
Clone: Isotype: Purification:	synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp. RB40716 IgG
Clone: Isotype: Purification: Target Details	synthetic peptide between 403-431 amino acids from the C-terminal region of mouse Lrmp. RB40716 IgG This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Molecular Weight:	59588
NCBI Accession:	NP_001268909
UniProt:	Q60664


Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Images



Western Blotting

Image 1. Western blot analysis of lysate from mouse spleen tissue lysate, using Mouse Lrmp Antibody (C-term) (ABIN1881507 and ABIN2838781). (ABIN1881507 and ABIN2838781) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 μg per lane.