

Datasheet for ABIN1881605  
**anti-OR52J3 antibody (C-Term)**[Go to Product page](#)**1** Image**1** Publication

## Overview

Quantity:	400 µL
Target:	OR52J3
Binding Specificity:	AA 283-311, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR52J3 antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	This OR52J3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 283-311 amino acids from the C-terminal region of human OR52J3.
Clone:	RB40815
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	OR52J3
Alternative Name:	OR52J3 ( <a href="#">OR52J3 Products</a> )
Background:	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response

## Target Details

that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

Molecular Weight: 35091

NCBI Accession: [NP\\_001001916](#)

UniProt: [Q8NH60](#)

## Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

## Publications

Product cited in: Hyrskyluoto, Bruelle, Lundh, Do, Kivinen, Rappou, Reijonen, Waltimo, Petersén, Lindholm, Korhonen: "Ubiquitin-specific protease-14 reduces cellular aggregates and protects against mutant huntingtin-induced cell degeneration: involvement of the proteasome and ER stress-activated kinase IRE1?." in: **Human molecular genetics**, Vol. 23, Issue 22, pp. 5928-39, (2014) ([PubMed](#)).


Davila, Froeling, Tan, Bonnard, Boland, Snippe, Hibberd, Seielstad: "New genetic associations detected in a host response study to hepatitis B vaccine." in: **Genes and immunity**, Vol. 11, Issue 3, pp. 232-8, (2010) ([PubMed](#)).

Chen, Qin, Li, Walters, Wilson, Mei, Wilson: "The proteasome-associated deubiquitinating enzyme Usp14 is essential for the maintenance of synaptic ubiquitin levels and the development of neuromuscular junctions." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 29, Issue 35, pp. 10909-19, (2009) ([PubMed](#)).

Nagai, Kadowaki, Maruyama, Takeda, Nishitoh, Ichijo: "USP14 inhibits ER-associated degradation via interaction with IRE1alpha." in: **Biochemical and biophysical research communications**, Vol. 379, Issue 4, pp. 995-1000, (2009) ([PubMed](#)).

Mines, Goodwin, Limbird, Cui, Fan: "Deubiquitination of CXCR4 by USP14 is critical for both CXCL12-induced CXCR4 degradation and chemotaxis but not ERK activation." in: **The Journal of biological chemistry**, Vol. 284, Issue 9, pp. 5742-52, (2009) ([PubMed](#)).

## Images



### Western Blotting

**Image 1.** Anti-OR52J3 Antibody (C-term) at 1:1000 dilution + whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.