

Datasheet for ABIN1881653
anti-PIGB antibody (N-Term)[Go to Product page](#)

1 Image

4 Publications

Overview

Quantity:	400 µL
Target:	PIGB
Binding Specificity:	AA 28-56, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIGB antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This PIGB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 28-56 amino acids from the N-terminal region of human PIGB.
Clone:	RB32518
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	PIGB
Alternative Name:	PIGB (PIGB Products)
Background:	This gene encodes a transmembrane protein that is located in the endoplasmic reticulum and

Target Details

is involved in GPI-anchor biosynthesis. The glycosylphosphatidylinositol (GPI) anchor is a glycolipid found on many blood cells and serves to anchor proteins to the cell surface. This gene is thought to encode a member of a family of dolichol-phosphate-mannose (Dol-P-Man) dependent mannosyltransferases.

Molecular Weight: 65056

NCBI Accession: [NP_004846](#)

UniProt: [Q92521](#)

Pathways: [Inositol Metabolic Process](#)

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Expiry Date: 6 months

Publications

Product cited in: Hyrskyluoto, Bruelle, Lundh, Do, Kivinen, Rappou, Reijonen, Waltimo, Petersén, Lindholm, Korhonen: "Ubiquitin-specific protease-14 reduces cellular aggregates and protects against mutant huntingtin-induced cell degeneration: involvement of the proteasome and ER stress-activated kinase IRE1?". in: **Human molecular genetics**, Vol. 23, Issue 22, pp. 5928-39, (2014) ([PubMed](#)).

Davila, Froeling, Tan, Bonnard, Boland, Snippe, Hibberd, Seielstad: "New genetic associations detected in a host response study to hepatitis B vaccine." in: **Genes and immunity**, Vol. 11,


Issue 3, pp. 232-8, (2010) ([PubMed](#)).

Chen, Qin, Li, Walters, Wilson, Mei, Wilson: "The proteasome-associated deubiquitinating enzyme Usp14 is essential for the maintenance of synaptic ubiquitin levels and the development of neuromuscular junctions." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 29, Issue 35, pp. 10909-19, (2009) ([PubMed](#)).

Nagai, Kadowaki, Maruyama, Takeda, Nishitoh, Ichijo: "USP14 inhibits ER-associated degradation via interaction with IRE1alpha." in: **Biochemical and biophysical research communications**, Vol. 379, Issue 4, pp. 995-1000, (2009) ([PubMed](#)).

Mines, Goodwin, Limbird, Cui, Fan: "Deubiquitination of CXCR4 by USP14 is critical for both CXCL12-induced CXCR4 degradation and chemotaxis but not ERK activation." in: **The Journal of biological chemistry**, Vol. 284, Issue 9, pp. 5742-52, (2009) ([PubMed](#)).

Images



Western Blotting

Image 1. All lanes : Anti-PIGB Antibody (N-term) at 1:1000 dilution Lane 1: Hela whole cell lysate Lane 2: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.