antibodies -online.com





anti-TSC1 antibody (pSer1130)



2

Publications



Go to Product page

_					
	W	0	rv	10	W

Quantity:	400 μL	
Target:	TSC1	
Binding Specificity:	pSer1130	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TSC1 antibody is un-conjugated	
Application:	Dot Blot (DB)	
Draduat Dataila		
Product Details		
Immunogen:	This mouse TSC1 Antibody is generated from rabbits immunized with a KLH conjugated	
	This mouse TSC1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse	
	synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse	
Immunogen:	synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse TSC1.	
Immunogen: Clone:	synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse TSC1. RB41227	
Immunogen: Clone: Isotype:	synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse TSC1. RB41227 Ig Fraction	
Immunogen: Clone: Isotype: Purification:	synthetic phosphopeptide corresponding to amino acid residues surrounding S1130 of mouse TSC1. RB41227 Ig Fraction	

Target Details

Background:	In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling (By similarity). Implicated as a tumor suppressor. Involved in microtubule-mediated protein	
	transport, but this seems to be due to unregulated mTOR signaling (By similarity).	
Molecular Weight:	128746	
NCBI Accession:	NP_075025	
UniProt:	Q9EP53	
Pathways:	RTK Signaling, AMPK Signaling, Regulation of Cell Size, Tube Formation	
Application Details		
Application Notes:	DB: 1:500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	
Publications		
Product cited in:	Mehta, Vazquez, Kulkarni, Kerrigan, Atwal, Metsugi, Toppmeyer, Levine, Hirshfield: "Polymorphic	
Guact cited iiii	Menta, Vazquez, Kulkarni, Kerngan, Atwai, Metsugi, Toppmeyer, Levine, Misimela. Folymorphic	
.,	variants in TSC1 and TSC2 and their association with breast cancer phenotypes." in: Breast	


Hoogeveen-Westerveld, Exalto, Maat-Kievit, van den Ouweland, Halley, Nellist: "Analysis of TSC1 truncations defines regions involved in TSC1 stability, aggregation and interaction." in: **Biochimica et biophysica acta**, Vol. 1802, Issue 9, pp. 774-81, (2010) (PubMed).

Mieulet, Lamb: "Tuberous sclerosis complex: linking cancer to metabolism." in: **Trends in molecular medicine**, Vol. 16, Issue 7, pp. 329-35, (2010) (PubMed).

Guo, Ying, Zhang, Yuan, Qian, Wang, Yang, He: "Tandem affinity purification and identification of the human TSC1 protein complex." in: **Acta biochimica et biophysica Sinica**, Vol. 42, Issue 4, pp. 266-73, (2010) (PubMed).

Liu, Wu, Chen, Ter-Minassian, Asomaning, Zhai, Wang, Su, Heist, Kulke, Lin, Liu, Christiani: "A Large-scale genetic association study of esophageal adenocarcinoma risk." in: **Carcinogenesis**, Vol. 31, Issue 7, pp. 1259-63, (2010) (PubMed).

Images



Dot Blot

Image 1. Dot blot analysis of mouse TSC1 Antibody (Phospho) Phospho-specific Pab (ABIN1881911 and ABIN2839922) on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6 µg per ml.