antibodies -online.com





anti-USP5 antibody (N-Term)

2 Images



Publications



Go to Product page

Overview	
Quantity:	400 μL
Target:	USP5
Binding Specificity:	AA 59-90, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	This USP5 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 59-90 amino acids from the N-terminal region of human USP5.
Clone:	RB4271
Isotype:	lg Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
	dialysis against PBS.
Target Details	
Target:	USP5
Alternative Name:	USP5 (USP5 Products)

Target Details

Background:

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),1 OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

Molecular Weight:

95786

NCBI Accession:

NP_001092006, NP_003472

UniProt:

P45974

Application Details

Application Notes:

WB: 1:1000. IHC-P: 1:50~100

Restrictions:

For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

Product cited in:

Mishra, Chandravanshi, Trigun, Krishnamurthy: "Ambroxol modulates 6-Hydroxydopamine-induced temporal reduction in Glucocerebrosidase (GCase) enzymatic activity and Parkinson's disease symptoms." in: **Biochemical pharmacology**, Vol. 155, pp. 479-493, (2019) (PubMed).

Images



A2058 130 95 - ◀ 72 55 -36 28

Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.

Western Blotting

Image 2. Western blot analysis of anti-USP5 Antibody (Nterm) (ABIN1882150 and ABIN2839176) in cell line lysates (35 μ g/lane). USP5(arrow) was detected using the purified Pab.