

Datasheet for ABIN1882166
anti-DAPK2 antibody (N-Term)

3 Images

6 Publications

[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	DAPK2
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAPK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This DAPK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human DAPK2.
Clone:	RB3053
Isotype:	Ig Fraction
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Target Details

Target:	DAPK2
Alternative Name:	DAPK2 (DAPK2 Products)

Target Details

Background: DAPK2 belongs to the serine/threonine protein kinase family. This protein contains a N-terminal protein kinase domain followed by a conserved calmodulin-binding domain with significant similarity to that of death-associated protein kinase 1 (DAPK1), a positive regulator of programmed cell death. Overexpression of this gene was shown to induce cell apoptosis. It uses multiple polyadenylation sites.

Molecular Weight: 42898

NCBI Accession: [NP_055141](#)

UniProt: [Q9UIK4](#)

Application Details

Application Notes: WB: 1:1000. WB: 1:2000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications


Product cited in: Tekin, Erden, Ozyalin, Cigremis, Colak, Sandal: "The effects of intracerebroventricular infusion of irisin on feeding behaviour in rats." in: **Neuroscience letters**, Vol. 645, pp. 25-32, (2017) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.



Western Blotting

Image 2. Western blot analysis of lysates from SH-SY5Y cell line and mouse brain tissue lysates (from left to right), using DK2 Antibody (M1) 7033a. 7033a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.



Western Blotting

Image 3. Anti-DK2 Antibody (M1) at 1:2000 dilution + HeLa whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.