

Datasheet for ABIN1882168
anti-GSG2 antibody (AA 323-352)



[Go to Product page](#)

2 Images

3 Publications

Overview

Quantity:	400 µL
Target:	GSG2
Binding Specificity:	AA 323-352
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GSG2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This Haspin GSG2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 323~352 amino acids from the central region of human haspin.
Clone:	RB3404
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	GSG2
Alternative Name:	Haspin GSG2 (GSG2 Products)
Background:	Post-translational modifications of conserved N-terminal tail residues in histones regulate

Target Details

many aspects of chromosome activity. Mitotic phosphorylation of H3 Thr 3 occurs in prophase and dephosphorylation during anaphase. Haspin, a dual serine/threonine kinase, plays an important role in regulation of chromosome and spindle function during mitosis and meiosis via its function in phosphorylation of the threonine residue in the third position of histone 3 (Thr3).

Molecular Weight: 88495

NCBI Accession: [NP_114171](#)

UniProt: [Q8TF76](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

Publications

Product cited in: Dai, Sultan, Taylor, Higgins: "The kinase haspin is required for mitotic histone H3 Thr 3 phosphorylation and normal metaphase chromosome alignment." in: **Genes & development**, Vol. 19, Issue 4, pp. 472-88, (2005) ([PubMed](#)).


Higgins: "The Haspin gene: location in an intron of the integrin alphaE gene, associated transcription of an integrin alphaE-derived RNA and expression in diploid as well as haploid cells." in: **Gene**, Vol. 267, Issue 1, pp. 55-69, (2001) ([PubMed](#)).

Tanaka, Iguchi, Nakamura, Kohroki, de Carvalho, Nishimune: "Cloning and characterization of human haspin gene encoding haploid germ cell-specific nuclear protein kinase." in: **Molecular human reproduction**, Vol. 7, Issue 3, pp. 211-8, (2001) ([PubMed](#)).



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated. BC = breast carcinoma, HC = hepatocarcinoma.



Western Blotting

Image 2. All lanes : Anti-GSG2 Antibody (Center) at 1:1000 dilution Lane 1: human testis lysate Lane 2: Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 88 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.