

## Datasheet for ABIN1885631 anti-APEX1 antibody (AA 1-15)



## Overview

| Overview             |  |
|----------------------|--|
| Quantity:            | 100 μL   |
| Target:              | APEX1  |
| Binding Specificity: | AA 1-15  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This APEX1 antibody is un-conjugated   |
| Application:         | Western Blotting (WB), Immunofluorescence (IF)   |
| Product Details      |  |
| Immunogen:           | Synthetic peptide contain a sequence corresponding to a region within amino acids 1 and 15 of  |
|                      | APE1   |
| Purification:        | Affinity purified by Protein A.  |
| Target Details       |  |
| Target:              | APEX1  |
| Alternative Name:    | APE1 (APEX1 Products)  |
| Background:          | Apurinic/apyrimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis,  |
|                      | by DNA damaging agents or by DNA glycosylases that remove specific abnormal bases.AP           |
|                      | sites are pre-mutagenic lesions that can prevent normal DNA replication so the cell contains   |
|                      | systems to identify and repair such sites. Class II AP endonucleases cleave the phosphodiester |

## **Target Details**

Storage Comment:

| l arget Details     |  |
|---------------------|--|
|                     | backbone 5' to the AP site. This gene encodes the major AP endonuclease in human cells. Splice |
|                     | variants have been found for this gene, all encode the same protein.[provided by RefSeq]       |
|                     | Synonyms: APE, APE-1, APE1, APEN, APEX, APX, HAP1, REF-1, REF1                                 |
| Molecular Weight:   | 36 kDa   |
| Gene ID:            | 328  |
| NCBI Accession:     | NP_001632, NM_001641   |
| Pathways:           | DNA Damage Repair, Chromatin Binding, Cell RedoxHomeostasis, Smooth Muscle Cell                |
|                     | Migration, Positive Regulation of Response to DNA Damage Stimulus                              |
| Application Details |  |
| Application Notes:  | Suggested dilutions:   |
|                     | Western blotting: 1.500-1.3000   |
|                     | Immunofluorescence: 1.100-1.200  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Buffer:             | 0.1 M Tris-buffered saline with 10 % Glycerol (pH 7.0).0.01 % Thimerosal was added as a        |
|                     | preservative.  |
| Preservative:       | Thimerosal (Merthiolate)   |
| Precaution of Use:  | Biohazard Informations: This product contains thimerosal which is hazardous.                   |
| Storage:            | 4 °C/-20 °C  |
|                     |  |

Store at -20 °C for long term preservation (recommended). Store at 4 °C for short term use.