

Datasheet for ABIN1885854
anti-IDH3G antibody (AA 1-257)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IDH3G
Binding Specificity:	AA 1-257
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IDH3G antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 1 and 257 of Human IDH3G
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	IDH3G
Alternative Name:	IDH3G (IDH3G Products)
Background:	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the

Target Details

mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. The protein encoded by this gene is the gamma subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase. This gene is a candidate gene for periventricular heterotopia. Several alternatively spliced transcript variants of this gene have been described, but only some of their full length natures have been determined. [provided by RefSeq]

Molecular Weight: 43 kDa

Gene ID: 3421

Application Details

Application Notes: Suggested dilutions:
Western blotting: 1.500-1.3000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 0.1 M Tris-buffered saline with 20 % Glycerol (pH 7.0). 0.01 % Thimerosal was added as a preservative.

Preservative: Thimerosal (Merthiolate)

Precaution of Use: Biohazard Informations: This product contains thimerosal which is hazardous.

Storage: 4 °C/-20 °C

Storage Comment: Store at -20 °C for long term preservation (recommended). Store at 4 °C for short term use.