antibodies -online.com





anti-HLA-DR Beta antibody (AA 1-262)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Overview		
Quantity:	100 μL	
Target:	HLA-DR Beta (HLA-DRB)	
Binding Specificity:	AA 1-262	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HLA-DR Beta antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	Recombinant protein fragment contain a sequence corresponding to a region within amino	
	acids 1 and 262 of Human HLA-DRB4	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	HLA-DR Beta (HLA-DRB)	
Abstract:	HLA-DRB Products	
Background:	HLA-DRB4 belongs to the HLA class II beta chain paralogues. This class II molecule is a	
	heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the	
	membrane.It plays a central role in the immune system by presenting peptides derived from	
	extracellular proteins.Class II molecules are expressed in antigen presenting cells (APC: B	
heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchore membrane. It plays a central role in the immune system by presenting peptides		

lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogues DRB3, DRB4 and DRB5. The presence of DRB4 is linked with allelic variants of DRB1, otherwise it is omitted. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9. [provided by RefSeq]

Molecular Weight: 30 kDa

Gene ID: 3126

NCBI Accession: NM_021983, NP_068818

Application Details

Application Notes: Suggested dilutions:

Western blotting: 1.500-1.3000

Restrictions: For Research Use only

Handling

Format:	Liquid	
Buffer:	0.1 M Tris-buffered saline with 10 % Glycerol (pH 7.0).0.01 % Thimerosal was added as a preservative.	
Preservative:	Thimerosal (Merthiolate)	
Precaution of Use:	Biohazard Informations: This product contains thimerosal which is hazardous.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at -20 °C for long term preservation (recommended). Store at 4 °C for short term use.	