



[Go to Product page](#)

Datasheet for ABIN1886504 anti-FBXL6 antibody (AA 478-539)

Overview

Quantity:	100 µL
Target:	FBXL6
Binding Specificity:	AA 478-539
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXL6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC)

Product Details

Immunogen:	Synthetic peptide contain a sequence corresponding to a region within amino acids 478 and 539 of Human FBXL6
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	FBXL6
Abstract:	FBXL6 Products
Background:	This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws

Target Details

containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class and, in addition to an F-box, contains several tandem leucine-rich repeats. Alternative splicing of this gene generates 2 transcript variants. [provided by RefSeq]

Molecular Weight: 59 kDa

Gene ID: 26233

NCBI Accession: [NM_012162](#), [NP_036294](#)

Application Details

Application Notes: Suggested dilutions:
Western blotting: 1.500-1.3000
Immunohistochemistry: 1.100-1.250
Immunofluorescence: 1.100-1.200

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 0.1 M Tris-buffered saline with 10 % Glycerol (pH 7.0). 0.01 % Thimerosal was added as a preservative.

Preservative: Thimerosal (Merthiolate)

Precaution of Use: Biohazard Informations: This product contains thimerosal which is hazardous.

Storage: 4 °C/-20 °C

Storage Comment: Store at -20 °C for long term preservation (recommended). Store at 4 °C for short term use.