



Datasheet for ABIN1895988
anti-FKBP8 antibody (AA 42-276)



[Go to Product page](#)

1 Image

Overview

Quantity:	50 µL
Target:	FKBP8
Binding Specificity:	AA 42-276
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FKBP8 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 42 and 276 of FKBP8. Percent identity by BLAST analysis: Human (100%), Mouse, Rat (92%). Type of Immunogen: Recombinant protein
Isotype:	IgG
Specificity:	Human FKBP38 / FKBP8
Purification:	Immunoaffinity purified

Target Details


Target:	FKBP8
Alternative Name:	FKBP38 / FKBP8 (FKBP8 Products)
Background:	Name/Gene ID: FKBP8 Synonyms: FKBP8, 38 kDa FKBP, 38 kDa FK506-binding protein, FKBP-8, FK506-binding protein 8, FKBP38, HFKBP38, FKBP-38, Rotamase, FK506 binding protein 8, 38 kDa, FK506-binding protein 8 (38kD), FKBP38, PPlase FKBP8
Gene ID:	23770
Pathways:	Autophagy

Application Details

Application Notes:	Approved: IHC, IHC-P (7.5 µg/mL) Usage: Antibody reactive against recombinant protein, detects immunogen in Western blot.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.1 M Tris-glycine, pH 7.0, 10 % glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.



Immunohistochemistry

Image 1. Anti-FKBP38 / FKBP8 antibody IHC staining of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 7.5 ug/ml.