antibodies - online.com







anti-ZDHHC8 antibody (Internal Region)



Image



Overview

Quantity:	100 μg
Target:	ZDHHC8
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This ZDHHC8 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	ZDHHC8
Immunogen:	Peptide with sequence C-QRDHPQLKTPPSK, from the internal region of the protein sequence according to NP_037505.1.
Sequence:	QRDHPQLKTP PSK
Isotype:	IgG
Specificity:	This antibody is expected to recognise isoform 2 (NP_037505.1) only.
Cross-Reactivity:	Dog, Human, Mouse, Pig, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

rarget betails	
Target:	ZDHHC8
Alternative Name:	ZDHHC8 (ZDHHC8 Products)
Background:	ZDHHC8, zinc finger, DHHC-type containing 8 , ZDHHCL1, ZNF378 , membrane-associated DHHC8 zinc finger protein, zinc finger, DHHC domain containing 8, zinc finger, DHHC domain like containing 1
Gene ID:	29801, 27801, 303796
NCBI Accession:	NP_037505
Application Details	
Application Notes:	Western Blot: Approx 80-85 kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 81.4 kDa according to NP_037505.1). Recommended concentration: 1-3 μ g/mL. An additional band of 48 kDa was consistently observed, however this band was no Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerate at 4°C for a few weeks and still remain viable.




Image 1. ABIN190853 (0.1 ug/ml) staining of Human Brain (Frontal Cortex) lysate (35 ug protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.