



[Go to Product page](#)

Datasheet for ABIN1928310  
**anti-PVRL2 antibody (AA 270-499)**

1 Image

Overview

Quantity:	100 µL
Target:	PVRL2
Binding Specificity:	AA 270-499
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PVRL2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 270 and 499 (Q92692) of Nectin 2.  Type of Immunogen: Recombinant protein
Isotype:	IgG
Specificity:	Human PVRL2 / CD112
Purification:	Immunoaffinity purified

## Target Details

---

Target:	PVRL2
Alternative Name:	PVRL2 / CD112 ( <a href="#">PVRL2 Products</a> )
Background:	Name/Gene ID: PVRL2  Synonyms: PVRL2, HVEB, Herpesvirus entry protein B, Poliovirus receptor-like 2, Nectin-2, PRR2, PVRR2, CD112, CD112 antigen, Herpes virus entry mediator B, Herpesvirus entry mediator B
Gene ID:	5819
Pathways:	<a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Cell-Cell Junction Organization</a>

## Application Details


---

Application Notes:	Approved: IHC, IHC-P (7.5 µg/mL), WB (1:500 - 1:3000)
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.1 M Tris-glycine, pH 7.0, 10 % glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.



### Immunohistochemistry

**Image 1.** Anti-PVRL2 / CD112 antibody IHC staining of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 7.5 ug/ml.