



[Go to Product page](#)

Datasheet for ABIN1944729

anti-C13orf30 antibody (C-Term)

3 Images

1 Publication

Overview

Quantity:	400 µL
Target:	C13orf30
Binding Specificity:	AA 106-139, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C13orf30 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This C13orf30 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 106-139 amino acids from the C-terminal region of human C13orf30.
Clone:	RB46016
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	C13orf30
Alternative Name:	C13orf30 (C13orf30 Products)
Molecular Weight:	16362

Target Details

Gene ID: 144809

UniProt: [Q8N7L0](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:25. IHC-P: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

Publications

Product cited in: Huang, Wang, Xu, Lu, Xu, Li, Zhou, Sha: "Expression of a novel RAD23B mRNA splice variant in the human testis." in: **Journal of andrology**, Vol. 25, Issue 3, pp. 363-8, (2004) ([PubMed](#)).


Humphray, Oliver, Hunt, Plumb, Loveland, Howe, Andrews, Searle, Hunt, Scott, Jones, Ainscough, Almeida, Ambrose, Ashwell, Babbage, Babbage, Bagguley, Bailey, Banerjee, Barker, Barlow, Bates, Beasley et al.: "DNA sequence and analysis of human chromosome 9. ..." in: **Nature**, Vol. 429, Issue 6990, pp. 369-74, (2004) ([PubMed](#)).

Masutani, Sugasawa, Yanagisawa, Sonoyama, Ui, Enomoto, Takio, Tanaka, van der Spek, Bootsma: "Purification and cloning of a nucleotide excision repair complex involving the xeroderma pigmentosum group C protein and a human homologue of yeast RAD23." in: **The EMBO journal**, Vol. 13, Issue 8, pp. 1831-43, (1994) ([PubMed](#)).




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffin-embedded H.thyroid section using C13orf30 Antibody (C-term) (ABIN1944729 and ABIN2838567). (ABIN1944729 and ABIN2838567) was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Western Blotting

Image 2. Western blot analysis of lysates from MCF-7, SK-BR-3, B-3 cell line (from left to right), using C13orf30 Antibody (C-term) (ABIN1944729 and ABIN2838567). (ABIN1944729 and ABIN2838567) was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Immunohistochemical analysis of paraffin-embedded H.brain section using C13orf30 Antibody (C-term) (ABIN1944729 and ABIN2838567). (ABIN1944729 and ABIN2838567) was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.