antibodies - online.com







anti-PAR1 antibody

Image

Molecular Weight:

Gene ID:

UniProt:

development.

47441 Da

2149

P25116

Publications



0	1 /	-	K	/1	-	1 A
u	\/	\vdash	I \	/ I	\vdash	1/1

Quantity:	100 μg	
Target:	PAR1 (F2R)	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PAR1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Isotype:	lgG1	
Target Details		
Target:	PAR1 (F2R)	
Alternative Name:	Thrombin Receptor (F2R Products)	
Background:	High affinity receptor for activated thrombin coupled to G proteins that stimulate	

phosphoinositide hydrolysis. May play a role in platelets activation and in vascular

Target Details

Pathways:

Nuclear Receptor Transcription Pathway, Skeletal Muscle Fiber Development, Positive Regulation of Endopeptidase Activity, Protein targeting to Nucleus

Application Details

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02 %
	sodium azide and 50 % glycerol.

Preservative: Sodium azide

Precaution of Use:

WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Storage:

4 °C,-20 °C

Publications

Product cited in:


Huang, Wang, Xu, Lu, Xu, Li, Zhou, Sha: "Expression of a novel RAD23B mRNA splice variant in the human testis." in: **Journal of andrology**, Vol. 25, Issue 3, pp. 363-8, (2004) (PubMed).

Humphray, Oliver, Hunt, Plumb, Loveland, Howe, Andrews, Searle, Hunt, Scott, Jones, Ainscough, Almeida, Ambrose, Ashwell, Babbage, Babbage, Bagguley, Bailey, Banerjee, Barker, Barlow, Bates, Beasley et al.: "DNA sequence and analysis of human chromosome 9. ..." in:

Nature, Vol. 429, Issue 6990, pp. 369-74, (2004) (PubMed).

Masutani, Sugasawa, Yanagisawa, Sonoyama, Ui, Enomoto, Takio, Tanaka, van der Spek, Bootsma: "Purification and cloning of a nucleotide excision repair complex involving the xeroderma pigmentosum group C protein and a human homologue of yeast RAD23." in: **The EMBO journal**, Vol. 13, Issue 8, pp. 1831-43, (1994) (PubMed).

Images



Western Blotting

Image 1. Western blot analysis of lysates from K562 cell line ,using Thrombin Receptor Antibody (ABIN484035 and ABIN1533416). ABIN484035 and ABIN1533416 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysates at 35 μ g.