antibodies .- online.com





anti-JARID2 antibody (N-Term)

2 Images




Go to Product page

Overview

Quantity:	0.1 mL
Target:	JARID2
Binding Specificity:	AA 1-100, N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This JARID2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF), Chromatin Immunoprecipitation (ChIP)
Product Details	
Product Details Immunogen:	A synthetic peptide made to an N-terminal portion of the human JARID2 or jumonji protein sequence (AA 1-100).
Immunogen:	sequence (AA 1-100).


Target Details

Target Details	
Target:	JARID2
Alternative Name:	JARID2 / Jumonji (JARID2 Products)
Pathways:	Chromatin Binding
Application Details	
Application Notes:	Chromatin Immunoprecipitation (ChIP), Immunocytochemistry 1:50-1:200,
	Immunofluorescence 1:50-1:200 , Western Blot 2 μ g/mL, Dilutions listed as a recommendation.
	Optimal dilution should be determined by investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1.83 mg/mL
Buffer:	Tris-glycine 150mM NaCl with 0.05% Sodium Azide as a preservative.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Repeated freeze/thaw cycles
Storage:	-20 °C
Storage Comment:	Store frozen. Aliquot as undiluted antisera and immediately place at -20°C. Antisera may have
	become trapped in top of vial during shipping. Centrifugation of vial is recommended before
	opening. Stable for at least 6 months at -20°C. Compromise the integrity of the antiserum.



Immunofluorescence

Image 1.



Western Blotting

Image 2.