antibodies -online.com





anti-SYNCRIP antibody (Biotin)





|)\/(| | | |
|------|--|--|--|
| | | | |

| Quantity: | 100 μg |
|--------------|--|
| Target: | SYNCRIP |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SYNCRIP antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)) |

Alternative Name:


Background:

| Product Details | |
|-----------------|--|
| Immunogen: | Recombinant human Heterogeneous nuclear ribonucleoprotein Q protein. |
| | |
| | Type of Immunogen: Recombinant protein |
| Isotype: | IgG |
| Specificity: | Human SYNCRIP / HnRNP Q |
| Purification: | Caprylic acid and ammonium sulfate precipitation |
| | |
| Target Details | |
| Target: | SYNCRIP |

SYNCRIP / HnRNP Q (SYNCRIP Products)

Name/Gene ID: SYNCRIP

| | Synonyms: SYNCRIP, DJ3J17.2, GRYRBP, HnRNP-Q, HNRPQ, HNRPQ1, HnRNP Q, NS1- |
|---------------------|--|
| | associated protein 1, NSAP1, GRY-RBP, PP68 |
| Gene ID: | 10492 |
| Application Dataile | |
| Application Details | |
| Application Notes: | Approved: ELISA, IHC, IHC-P, WB |
| | Usage: The applications listed have been tested for the unconjugated form of this product. |
| | Other forms have not been tested. |
| Comment: | Target Species of Antibody: Human |
| Restrictions: | For Research Use only |
| | |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, 0.03 % Proclin 300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be |
| | handled by trained staff only. |
| Handling Advice: | Avoid freeze-thaw cycles. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles. |



Immunohistochemistry

Image 1. Anti-SYNCRIP / HnRNP Q antibody IHC staining of human skin. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 20 ug/ml. This image was taken for the unconjugated form of thi ...