

Datasheet for ABIN1957173  
**anti-TNC antibody (FITC)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µg
Target:	TNC
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNC antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	Recombinant human Tenascin protein.
	Type of Immunogen: Recombinant protein
Isotype:	IgG
Specificity:	Human TNC / Tenascin C
Purification:	Caprylic acid and ammonium sulfate precipitation

## Target Details

Target:	TNC
Alternative Name:	TNC / Tenascin C ( <a href="#">TNC Products</a> )
Background:	Name/Gene ID: TNC

## Target Details


	Family: Fibronectin
	Synonyms: TNC, 150-225, Cytotactin, GP, Hexabrachion (tenascin), GMEM, GP 150-225, Miotendinous antigen, Myotendinous antigen, JI, Tenascin, Tenascin C, Neuronection, Tenascin-C isoform 14/AD1/16, TN-C, Hexabrachion, HXB, Tenascin-C, TN
Gene ID:	3371
Pathways:	<a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Regulation of Cell Size</a> , <a href="#">Skeletal Muscle Fiber Development</a>

## Application Details

Application Notes:	Approved: ELISA, IHC, IHC-P, WB
	Usage: The applications listed have been tested for the unconjugated form of this product. Other forms have not been tested.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, 0.03 % Proclin 300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid freeze-thaw cycles.
Storage:	-20 °C,-80 °C
Storage Comment:	Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles.



#### Immunohistochemistry

**Image 1.** Anti-TNC / Tenascin C antibody IHC staining of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 20 ug/ml. This image was taken for the unconjugated form of ...