

Datasheet for ABIN197234
anti-FAK antibody (Tyr861)



[Go to Product page](#)

2 Images

Overview

Quantity:	0.1 mL
Target:	FAK (PTK2)
Binding Specificity:	Tyr861
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAK antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	The antiserum was produced against synthesized non-phosphopeptide derived from human FAK around the phosphorylation site of tyrosine 861 (H-I-YP-Q-P).
Specificity:	This antibody detects endogenous levels of total FAK protein.
Purification:	Affinity Chromatography using epitope-specific phosphopeptide.

Target Details

Target:	FAK (PTK2)
Alternative Name:	FAK1 / PTK2 (PTK2 Products)
Background:	Focal adhesion kinase (FAK) is a non receptor protein tyrosine kinase discovered as a substrate for Src and as a key element of integrin signaling. FAK plays a central role in cell spreading, differentiation, migration, cell death and acceleration of the G1 to S phase transition of the cell

Target Details

cycle. FAK regulation includes phosphorylation at multiple tyrosine and serine residues. Phosphorylation of tyrosine generally is associated with positive regulation and growth promotion, however, dephosphorylation at these sites occurs as cells enter mitosis (M-Phase of the cell cycle). In contrast, serine phosphorylation either remains high or is increased as cells enter mitosis and may play a role in focal adhesion disassembly. FAK and its phosphorylation states have been implicated in cancer metastasis and tumor cell survival and adhesion-independent growth. Additionally, recent evidence indicates that elevation of FAK activity in human carcinoma cells is associated with increased invasive potential. A central role in tumor formation and progression suggests that FAK is an attractive target for therapeutic intervention. Synonyms: FAK1, FAK, Focal adhesion kinase 1, Protein-tyrosine kinase 2, pp125FAK

Gene ID:	5747
NCBI Accession:	NP_722560
UniProt:	Q05397
Pathways:	Response to Growth Hormone Stimulus , CXCR4-mediated Signaling Events , Smooth Muscle Cell Migration , Signaling of Hepatocyte Growth Factor Receptor , VEGF Signaling

Application Details

Application Notes:	Western Blot: 1/500approx. 1/1000. Immunohistochemistry on Paraffin Sections: 1/50approx. 1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only


Handling

Concentration:	1.0 mg/mL
Buffer:	PBS (without Mg2+ and Ca2+), pH 7.4 containing 150 mM NaCl, 0.02 % Sodium Azide and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.

Handling

Storage:	-20 °C
Storage Comment:	Store the antibody (in aliquots) at -20 °C.

Images



Peptide - +

Image 1.




Image 2.