



[Go to Product page](#)

Datasheet for ABIN1978852
anti-BCL2L10 antibody (AA 60-153)

1 Image

Overview

Quantity:	50 µL
Target:	BCL2L10
Binding Specificity:	AA 60-153
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic peptide corresponding to a region within amino acids 60 and 153 of BCL2L10 (SwissProt Q9HD36). Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Human BCL-B / Diva
Purification:	Immunoaffinity purified

Target Details

Target:	BCL2L10
---------	---------

Target Details


Alternative Name:	BCL2L10 / Diva (BCL2L10 Products)
Background:	Name/Gene ID: BCL2L10 Family: Apoptosis Synonyms: BCL2L10, BCL-B, Apoptosis regulator Bcl-B, BCLB, Bcl2-L-10, Diva not vaccine, Anti-apoptotic protein NrH, Bcl-2-like protein 10, Boo, Boo not bladder, Diva
Gene ID:	10017
Pathways:	Positive Regulation of Endopeptidase Activity

Application Details

Application Notes:	Approved: ICC (1:100 - 1:1000), IHC, IHC-P (7.5 µg/mL), WB (1:500 - 1:3000) Usage: IHC-paraffin: Suggested antigen retrieval using heat mediated 10 mM Citrate buffer (pH 6.0) or Tris-EDTA buffer (pH 8.0).
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.1 M Tris-glycine, pH 7.0, 10 % glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.



Immunohistochemistry

Image 1. Anti-BCL2L10 / Diva antibody IHC staining of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 7.5 ug/ml.