
# antibodies .- online.com





## anti-CIDEA antibody (AA 1-219)

2 Images



Go to Product page

#### Overview


Background:

Quantity:	100 μL
Target:	CIDEA
Binding Specificity:	AA 1-219
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	CIDEA (Met1-Gly219)
Immunogen:	CIDEA (Met1-Gly219) IgG
Isotype:	IgG  The antibody is a rabbit polyclonal antibody raised against CIDEA. It has been selected for its
Isotype: Specificity:	IgG  The antibody is a rabbit polyclonal antibody raised against CIDEA. It has been selected for its ability to recognize CIDEA in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification:	IgG  The antibody is a rabbit polyclonal antibody raised against CIDEA. It has been selected for its ability to recognize CIDEA in immunohistochemical staining and western blotting.

Alternative Names: CIDE-A, Cell death activator CIDE-A


### **Application Details**

Application Notes:	<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months
Images	



Western Blotting

Image 1.



#### **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded Brain tissue