

Datasheet for ABIN1980429

anti-Integrin alpha 1 antibody (AA 119-355)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	Integrin alpha 1 (ITGA1)
Binding Specificity:	AA 119-355
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Integrin alpha 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Integrin Alpha 1 (ITGa1)
Immunogen:	ITGa1 (Leu119-Leu355)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ITGa1. It has been selected for its ability to recognize ITGa1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	Integrin alpha 1 (ITGA1)
---------	--------------------------

Target Details


Alternative Name:	ITGa1 (ITGA1 Products)
Background:	CD49-A, CD49a, ITG-A1, VLA1, CD49 antigen-like family member A, Laminin and collagen receptor
Pathways:	EGFR Signaling Pathway , CXCR4-mediated Signaling Events , Signaling of Hepatocyte Growth Factor Receptor , Integrin Complex

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded Kidney tissue