

Datasheet for ABIN1981845

STAT6 ELISA Kit

5 Images

1 Publication

[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	STAT6
Binding Specificity:	pTyr641
Reactivity:	Human, Rat, Mouse
Method Type:	Cell ELISA
Application:	ELISA

Product Details

Purpose:	Cell-Based Human/Mouse Stat 6 (Tyr641) Phosphorylation ELISA Kit. Suitable for adherent whole cell lines.
Brand:	CellBIND®
Sample Type:	Cell Culture Cells
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibodies provided in this kit recognizes human and mouse Stat 6 phosphorylated at site Tyrosine-641 and total Stat 6 for comparison.
Characteristics:	<ul style="list-style-type: none">• Site and signal pathway-specific• In vitro detection of adherent cell culture• No sample lysis needed• Compatible with a standard ELISA plate reader• Faster results than with ELISA

Product Details

- Adaptable for high-throughput screening and drug discovery

Components:

- uncoated 96-well Microplate
- Wash Buffer A
- Wash Buffer B
- Fixing Solution
- Quenching Buffer
- Blocking Buffer
- Anti-phospho antibody
- Anti-pan antibody
- HRP-Conjugated Secondary Antibody
- TMB One-Step Substrate
- Stop Solution

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: STAT6

Alternative Name: STAT6 ([STAT6 Products](#))

Gene ID: 6778

UniProt: [P42226](#)

Pathways: [JAK-STAT Signaling](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#)

Application Details

Sample Volume: 100 µL

Plate: Uncoated

Protocol:

1. Seed 10,000-30,000 cells into each well and incubate overnight.
2. Apply various treatment, inhibitors or activators according to manufacture's instructions.

3. Add 100 µL of Fixing Solution into each well and incubate for 20 min at RT with shaking.
4. Add 200 µL of prepared 1X Quenching Buffer and incubate 20 min at RT.
5. Add 200 µL of Blocking Solution and incubate for 1 h at 37 °C.
6. Add 50 µL of 1X anti-phospho-protein specific antibody or anti-pan-protein specific antibody to each well and incubate for 2 h at RT.
7. Add 50 µL of prepared 1X HRP-Anti-Rabbit or Mouse IgG and incubate for 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

Reagent Preparation:	<p>NOTE: Thaw all reagents to room temperature immediately before use. If wash buffers contain visible crystals, warm to room temperature and mix gently until dissolved.</p> <p>NOTE: Briefly centrifuge (~1,000g) ITEMS G, H, and I before opening to ensure maximum recovery.</p> <p>For more information look at the picture.</p>
----------------------	---

Assay Procedure:	<p>NOTE: ALL incubations and wash steps must be performed under gentle rocking or rotation (~1-2 cycles/sec).</p> <p>1. Design your experiment. For example, see Figure 2 below.</p> <p>OPTIONAL: If seeding HUVECs, HMEC-1 or other loosely attached cells, coat the Uncoated 96-Well Microplate (ITEM A) by adding 100 µL poly-L-Lysine (Recommended Sigma Aldrich, Cat#: P4832) into each well and then follow manufacturer's instructions. A pre-coated CellBIND® microplate or other poly-lysine treated tissue culture plate may be used in place of Item A.</p> <p>2. Seed 100 µL of 30,000 cells into each well of the Uncoated 96-Well Microplate (ITEM A) provided and incubate overnight at 37 °C with 5 % CO₂.</p> <p>NOTE: The optimal cell number used will vary on the cell line and the relative amount of protein phosphorylation. More or less cells may be used but this must be determined empirically.</p> <p>NOTE: The cells can be starved ~4-24 hours (depending on cell line) prior to treatment with inhibitors or activators.</p> <p>3. Apply various treatments, inhibitors (such as siRNA or chemicals) or activators according to manufacturer's instructions and incubate for the desired time points.</p> <p>NOTE: It is recommended to dissolve inhibitors or activators into serum-free cell culture medium before treating the cells (unless otherwise stated in the manufacturer's instructions.)</p> <p>4. Discard the cell culture medium by flipping the microplate upside down and gently tapping the bottom of the microplate over a sink.</p> <p>5. Wash by pipetting 200 µL of the prepared 1X Wash Buffer A (ITEM B) into each well. Discard the wash buffer (same as step 4) and wash 2 more times for a total of 3 washes using fresh</p>
------------------	--

wash buffer each time. After the final wash, gently blot the microplate onto a paper towel to remove any excess/remaining buffer.

NOTE: To avoid cell loss, do not pipette directly onto the cells. Instead, gently dispense the liquid down the wall of cell culture wells. Also avoid the use of vacuum suction or too forcefully tapping the microplate when discarding any solution.

6. Add 100 μ L of Fixing Solution (ITEM D) into each well and incubate for 20 minutes at room temperature.

NOTE: The fixing solution is used to permeabilize the cells.

7. Repeat wash step 5.

8. Add 200 μ L of the prepared 1X Quenching Buffer (ITEM E) into each well and incubate 20 minutes at room temperature.

NOTE: The quenching buffer is used to minimize the background response.

9. Wash 4 times with 1X Wash Buffer A.

10. Add 200 μ L of the prepared 1X Blocking Buffer (ITEM F) into each well and incubate for 1 hour at 37 °C.

11. Wash 3 times with the prepared 1X Wash Buffer B (ITEM C).

NOTE: If needed, the microplate may be stored at -80 °C for several days after this wash.

12. Add 50 μ L of the prepared 1X primary antibody (ITEM G or H) into each corresponding well and incubate for 2 hours at room temperature.

13. Wash 4 times with 1X Wash Buffer B.

14. Add 50 μ L of the prepared 1X HRP Conjugated secondary antibody (ITEM I-1 or I-2) into each well and incubate for 1 hour at room temperature.

NOTE: Item I-1 is the secondary antibody for Item G (primary antibody). Item I-2 is the secondary antibody for Item H (primary antibody).

15. Wash 4 times with 1X Wash Buffer B.

16. Add 100 μ L of the TMB Substrate (ITEM J) into each well and incubate for 30 minutes at room temperature in the dark.

17. Add 50 μ L of the Stop Solution (ITEM K) into each well. Read at 450 nm immediately.

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
------------------	------------------------------------

Storage:	-20 °C
----------	--------

Storage Comment:	The entire kit may be stored at -20°C for up to 6 months from the date of shipment. Avoid
------------------	---

Handling

repeated freeze-thaw cycles.

Expiry Date: 6 months

Publications

Product cited in: Patel, Wang, Lee, Taylor, Pierce, LaRochelle: "Stat6 and Jak1 are common elements in platelet-derived growth factor and interleukin-4 signal transduction pathways in NIH 3T3 fibroblasts." in: **The Journal of biological chemistry**, Vol. 271, Issue 36, pp. 22175-82, (1996) ([PubMed](#)).

Images




Image 1. Example of how to seed cells for cell-based assay

	ITEM	COMPONENT	PREPARATION	EXAMPLE
	A	Uncoated 96-Well Microplate	No Preparation	N/A
	B	20X Wash Buffer A Concentrate	Dilute 20-fold with distilled or deionized water	25 ml of concentrate + 475 ml of water
	C	20X Wash Buffer B Concentrate		500 ml of 1X working solution
	D	Fixing Solution	No Preparation	N/A
	E	30X Quenching Buffer Concentrate	Dilute 30-fold with 1X Wash Buffer A	1 ml of concentrate + 29 ml of wash buffer = 30 ml of 1X working solution
	F	5X Blocking Buffer Concentrate	Dilute 5-fold with distilled or deionized water	20 ml of concentrate + 80 ml of water = 100 ml of 1X working solution
PRIMARY ANTIBODY	G	500X Rabbit Anti-phospho (Tyr641) STAT6 Concentrate	Dilute 500-fold with 1X Blocking Buffer	10 μ l of concentrate + 4990 μ l of 1X Blocking Buffer = 5 ml of 1X working solution
	H	5000X Mouse Anti-STAT6 Concentrate	Dilute 5000-fold with 1X Blocking Buffer	2 μ l of concentrate + 9998 μ l of 1X Blocking Buffer = 10 ml of 1X working solution
	I-1	1000X HRP Conjugated Anti-Rabbit IgG Concentrate		10 μ l of concentrate + 9990 μ l of 1X Blocking Buffer = 10 ml of 1X working solution
SECONDARY ANTIBODY	I-2	1000X HRP Conjugated Anti-Mouse IgG Concentrate	Dilute 1000-fold with 1X Blocking Buffer	
	J	TMB Substrate		
	K	Stop Solution	No Preparation	N/A

Image 2. This picture shows the reagent preparation.




Image 3. Cell-Based protein phosphorylation procedure

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN1981845.