

Datasheet for ABIN1981880

anti-CD79a antibody (AA 208-222) (APC)



[Go to Product page](#)

1 Image

7 Publications

Overview

Quantity:	100 tests
Target:	CD79a (CD79A)
Binding Specificity:	AA 208-222
Reactivity:	Human, Mouse, Rat, Cow, Pig, Chicken, Rabbit, Horse, Guinea Pig, Dog, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD79a antibody is conjugated to APC
Application:	Intracellular Flow Cytometry (ICFC)

Product Details

Purpose:	Anti-Hu CD79a APC
Immunogen:	Synthetic peptide corresponding to C terminal amino acids 208-222 of human CD79a
Clone:	HM47
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody HM47 reacts with intracellular domain of CD79a (Ig alpha), a 40-45 kDa subunit of B cell antigen-specific receptor (BCR) and its early developmental forms.
Cross-Reactivity (Details):	Human, Non-Human Primates, Porcine, Mouse, Rat, Bovine, Canine (Dog), Equine (Horse), Guinea pig, Rabbit, Chicken
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion

Product Details

chromatography.

Target Details

Target:	CD79a (CD79A)
Alternative Name:	CD79a (CD79A Products)
Background:	CD79a molecule,CD79a (Ig alpha, MB1) forms disulfide-linked heterodimer with CD79b (Ig beta). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. At the plasma cell stage, CD79a is present as an intracellular component. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.,BCR alpha, Ig-alpha, MB-1, IGA
Gene ID:	973
UniProt:	P11912
Pathways:	BCR Signaling

Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. Intracellular staining.
Restrictions:	For Research Use only

Handling

Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Handling Advice: Do not freeze. Avoid prolonged exposure to light.
Do not use after expiration date stamped on vial label.
Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.

Storage: 4 °C

Storage Comment: Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Publications

Product cited in: Zhao, Hassan, Perry, Ning, Stass, Dehner: "C-MYC rearrangements are frequent in aggressive mature B-Cell lymphoma with atypical morphology." in: **International journal of clinical and experimental pathology**, Vol. 1, Issue 1, pp. 65-74, (2008) ([PubMed](#)).


Bhargava, Kallakury, Ross, Azumi, Bagg: "CD79a is heterogeneously expressed in neoplastic and normal myeloid precursors and megakaryocytes in an antibody clone-dependent manner." in: **American journal of clinical pathology**, Vol. 128, Issue 2, pp. 306-13, (2007) ([PubMed](#)).

Fernandez, West, Jackson, Kidney: "Immunohistochemical and histochemical stains for differentiating canine cutaneous round cell tumors." in: **Veterinary pathology**, Vol. 42, Issue 4, pp. 437-45, (2005) ([PubMed](#)).

Islas-Ohlmayer, Padgett-Thomas, Domiati-Saad, Melkus, Cravens, Martin, Netto, Garcia: "Experimental infection of NOD/SCID mice reconstituted with human CD34+ cells with Epstein-Barr virus." in: **Journal of virology**, Vol. 78, Issue 24, pp. 13891-900, (2004) ([PubMed](#)).

Torlakovic, Torlakovic: "B-cell markers in lymphocyte predominance Hodgkin disease." in: **Archives of pathology & laboratory medicine**, Vol. 126, Issue 7, pp. 862-3, (2002) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)



Flow Cytometry

Image 1. Flow cytometry analysis (intracellular staining) of CD79a in human peripheral blood with anti-CD79a (HM47) APC.