

Datasheet for ABIN1993192

anti-AMH antibody (AA 450-552) (FITC)



Overview

| Quantity: | 100 μg |
|----------------------|--|
| Target: | AMH |
| Binding Specificity: | AA 450-552 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AMH antibody is conjugated to FITC |
| Application: | Western Blotting (WB), ELISA |
| Product Details | |
| Immunogen: | Recombinant Mouse Muellerian-inhibiting factor protein (450-552AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Purification: | >95%, Protein G purified |
| Target Details | |
| Target: | AMH |
| Alternative Name: | Amh (AMH Products) |
| Background: | Background: This glycoprotein, produced by the Sertoli cells of the testis, causes regression of |

the Muellerian duct. It is also able to inhibit the growth of tumors derived from tissues of

Target Details

Storage Comment:

| rarget Details | |
|---------------------|---|
| | Muellerian duct origin. |
| | Aliases: Amh antibody, Muellerian-inhibiting factor antibody, Anti-Muellerian hormone antibody, |
| | AMH antibody, Muellerian-inhibiting substance antibody, MIS antibody |
| UniProt: | P27106 |
| Pathways: | Negative Regulation of Hormone Secretion |
| Application Details | |
| Application Notes: | Optimal working concentration should be determinde by the investigator. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 |
| | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin TM : a POISONOUS AND HAZARDOUS SUBSTANCE, which should |
| | be handled by trained staff only. |
| Handling Advice: | Avoid repeated freeze. |
| Storage: | -20 °C,-80 °C |
| | |

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.