

Datasheet for ABIN199368

anti-KIR2DL1 antibody (FITC)



	۱۱ /	\cap	r\/	i,	\sim 1	Λ/	
C	V	ヒ	ΙV	ľ	こ	٧V	

Overview	
Quantity:	100 μg
Target:	KIR2DL1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This KIR2DL1 antibody is conjugated to FITC
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP), Flow Cytometry (FACS)
Product Details	
Immunogen:	Human natural killer (NK) cell lysate.
leatune:	laC1

Immunogen:	Human natural killer (NK) cell lysate.
Isotype:	lgG1
Specificity:	Recognizes KIR2D members of the killer cell immunoglobulin (Ig)-like receptor (KIR) family:
	CD158a, CD158b and P50.3. Specifically recognizes the long and short forms CD158a and
	CD158b (KIR2DL, KIR2DS1 and KIR2DS2 respectively) and also p50.3 (KIR2DS4). Reported to
	have functional activity, activating NK cell cytotoxicity via KIR2DS and inhibiting via KIR2DL
	forms.
Purification:	Protein G purified

Target Details

Target:	KIR2DL1
Alternative Name:	KIR2DL1 / CD158a (KIR2DL1 Products)

Target Details

Background:	Name/Gene ID: KIR2DL1 Family: Immunoglobulin
	Synonyms: KIR2DL1, 47.11, CI-42, MHC class I NK cell receptor, NKAT-1, p58 NK receptor CL-42/47.11, NKAT, NKAT1, p58.1, CD158A, CD158a antigen, Killer Ig receptor, KIR-K64, KIR221
Gene ID:	3802
UniProt:	P43626

Application Details

Application Notes:	Approved: ELISA, Flo, IP, WB
	Usage: Suitable for use in Flow Cytometry, ELISA. Immunoprecipitation and Western Blot. Flow
	Cytometry: Neat-1:5. Use 10 μ L of the optimal working dilution to label 1 x10 6 in 100 μ L. The
	applications listed have been tested for the unconjugated form of this product. Other forms
	have not been tested.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.2, 1.0 % BSA, 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Short term: 4°C Long term: Add glycerol (40-50%) -20°C.