.-online.com antibodies

Datasheet for ABIN2017858 FGF2 Protein (AA 10-155)



Οv		

000000000000000000000000000000000000000	
Quantity:	50 µg
Target:	FGF2
Protein Characteristics:	AA 10-155
Origin:	Cow
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Product Details	
Characteristics:	ED50 < 1.0 ng/mL, measured by a cell proliferation assay using 3T3 cells, corresponding to a
	specific activity of > 1 x 10^6 units/mg.
Purity:	> 95 % by SDS-PAGE analysis.
Endotoxin Level:	< 0.2 EU/µg, determined by LAL method.
Target Details	
Target:	FGF2
Alternative Name:	Fibroblast Growth Factor-Basic (FGF-Basic) (FGF2 Products)
Background:	Fibroblast Growth Factor-basic (FGF-basic), also known as FGF-2, is a pleiotropic cytokine and
	one of the prototypic members of the heparin-binding FGF family. Like other FGF family
	members, FGF-basic has the beta trefoil structure. In vivo, FGF-basic is produced by a variety of

cells, including cardiomycotes, fibroblasts, and vascular cells. FGF-basic regulates a variety of

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2017858 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details

	processes including cell proliferation, differentiation, survival, adhesion, motility, apoptosis, limb
	formation and wound healing. FGF-basic can be tumorigenic due to its role in angiogenesis and
	blood vessel remodeling. The angiogenic effects of FGF-basic can produce beneficial
	cardioprotection during acute heart injury.Recombinant bovine Fibroblast Growth Factor-basic
	(rbFGF-basic) produced in E. coli is a single non-glycosylated polypeptide chain containing 146
	amino acids. A fully biologically active molecule, rbFGF-basic has a molecular mass of 16.4 kDa
	analyzed by reducing SDS-PAGE.
	Synonyms: FGF-2, BFGF, FGFB, HBGF-2
Molecular Weight:	16.4 kDa, observed by reducing SDS-PAGE.
UniProt:	P03969
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin
	Signaling Pathway, C21-Steroid Hormone Metabolic Process, Inositol Metabolic Process,
	Glycosaminoglycan Metabolic Process, Protein targeting to Nucleus, S100 Proteins

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstituted in ddH20 at 100 µg/mL.
Buffer:	Lyophilized after extensive dialysis against PBS.
Storage:	-80 °C
Storage Comment:	Lyophilized recombinant bovine Fibroblast Growth Factor-basic (rbFGF-basic) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rbFGF-basic remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.
Expiry Date:	6 months