

Datasheet for ABIN2017951 **CCL14 Protein (AA 22-93)**



[Go to Product page](#)

Overview

Quantity:	25 µg
Target:	CCL14
Protein Characteristics:	AA 22-93
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active

Product Details

Characteristics:	ED50 < 25 µg/mL, measured by the FLIPR assay using CHO cells transfected with human CCR5, the receptor of human CCL14, corresponding to a specific activity of > 40 units/mg.
Purity:	> 95 % by SDS-PAGE analysis.
Endotoxin Level:	< 0.2 EU/µg, determined by LAL method.

Target Details

Target:	CCL14
Abstract:	CCL14 Products
Background:	HCC-1/CCL14 is a member of the chemokine family, which are small chemotactic proteins that regulate cell migration under inflammatory and steady state conditions. HCC-1 is expressed in epithelial and decidual cells and is unique among chemokines due to its high abundance in normal human plasma. HCC-1 can bind to chemokine receptors CCR1 and CCR5, however full

Target Details

length HCC-1 is a weak agonist of CCR1 and only becomes potent after removal of its eight N-terminal residues. Chemokine decoy receptor D6 can bind HCC-1 and promote its degradation as a means to regulate its level in vivo. Functionally HCC-1 promotes trophoblast migration by regulating extracellular matrix components as well as specific adhesion

molecules. Recombinant human Hemofiltrate CC Chemokine-1 (72 a.a.) (HCC-1)/CCL14 (rhHCC-1) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 72 amino acids. A fully biologically active molecule, rhHCC-1 has a molecular mass of 8.4 kDa analyzed by reducing SDS-PAGE.

Synonyms: NCC-2, SCYA14

Molecular Weight: 8.4 kDa, observed by reducing SDS-PAGE.

UniProt: [Q16627](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstituted in ddH₂O at 100 µg/mL.

Buffer: Lyophilized after extensive dialysis against PBS.

Storage: -80 °C

Storage Comment: Lyophilized recombinant human Hemofiltrate CC Chemokine-1 (72 a.a.) (HCC-1)/CCL14 (rhHCC-1) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhHCC-1 remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

Expiry Date: 6 months