antibodies -online.com





Datasheet for ABIN2018235

CCL7 Protein



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Overview		
Quantity:	1 mg	
Target:	CCL7	
Origin:	Mouse	
Source:	Escherichia coli (E. coli)	
Protein Type:	Recombinant	
Biological Activity:	Active	
Product Details		
Sequence:	QPDGPNASTC CYVKKQKIPK RNLKSYRRIT SSRCPWEAVI FKTKKGMEVC AEAHQKWVEE AIAYLDMKTP TPKP	
Characteristics:	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human monocytes is in a concentration range of 100-300 ng/mL.	
Purity:	> 95 % by SDS-PAGE and HPLC analyses.	
Sterility:	0.2 µm filtered	
Endotoxin Level:	< 1 EU/µg of rMuMCP-3/CCL7 as determined by LAL method.	
Target Details		
Target:	CCL7	
Alternative Name:	MCP-3/CCL7 (CCL7 Products)	
Background:	Monocyte Chemotactic Protein-3 (MCP3) and CCL7 are two monocyte chemotactic proteins. Both MCP3 and CCL7 are members of the CC family of chemokines and share 62 % and 71 %	

amino acid sequence identity, respectively, with MCP1. CCL7 also shares 58 % amino acid		
identity with MCP3. Similarly to other CC chemokines, all three MCP proteins are monocyte		
chemoattractants. In addition, the three MCPs can chemoattract activated NK cells as well as		
CD4+ and CD8+ T lymphocytes. All three cytokines have also been shown to attract eosinophils		
and induce histamine secretion from basophils.		
Synonyms: Monocyte Chemotactic Protein-3, CCL7, MARC		

Molecular Weight:

8.5 kDa, a single, non-glycosylated polypeptide chain containing 74 amino acids.

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.	
Buffer:	Lyophilized from a 0.2 µm filtered concentrated solution in 2x PBS, pH 7.4.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C.	