

# Datasheet for ABIN2018300

# Oncostatin M Protein (OSM) (AA 25-205)



## Overview

Overview	
Quantity:	50 μg
Target:	Oncostatin M (OSM)
Protein Characteristics:	AA 25-205
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Product Details	
Characteristics:	ED50 < 10 ng/mL, measured by a cell proliferation assay using Rat Embryo Brain cells,
	corresponding to a specific activity of > 1x 10^5 units/mg.
	AA 25-205, expressed with an N-terminal Met.
Purity:	> 95 % as analyzed by SDS-PAGE and HPLC.
Endotoxin Level:	< 0.2 EU/µg, determined by LAL method.
Target Details	
Target:	Oncostatin M (OSM)
Alternative Name:	Oncostatin M(OSM) (OSM Products)
Background:	Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfam

which also includes IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor,

cardiotrophin-1, and novel neurotropin-1. In vivo, OSM is secreted from activated T cells,

monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and shares a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor beta or LIF Receptor beta to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells. Recombinant Mouse Oncostatin M (rmOSM) produced in E. coli is a single non-glycosylated polypeptide chain containing 182 amino acids. A fully biologically active molecule, rmOSM has a molecular mass of 20.5 kDa analyzed by reducing SDS-PAGE.

Synonyms: OSM

Molecular Weight:

20.5 kDa, observed by reducing SDS-PAGE.

UniProt:

P53347

Pathways:

JAK-STAT Signaling, Negative Regulation of Hormone Secretion

## **Application Details**

Restrictions:

For Research Use only

## Handling

Format:	Lyophilized
Reconstitution:	Reconstituted in ddH2O or PBS at 100 μg/mL.
Buffer:	Lyophilized after extensive dialysis against PBS.
Storage:	-80 °C
Storage Comment:	Lyophilized recombinant Mouse Oncostatin M (rmOSM) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rmOSM remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.
Expiry Date:	6 months