



[Go to Product page](#)

Datasheet for ABIN2115487
NRG1-beta 1 Protein (AA 1-246) (His tag)

Overview

Quantity:	50 µg
Target:	NRG1-beta 1
Protein Characteristics:	AA 1-246
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NRG1-beta 1 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human Pro-Neuregulin-1/NRG1-β1/HRG1-β1 (Met1-Lys246, N-6His)
Sequence:	MGSSHHHHHH SSSLVPRGSH MSERKEGRGK GKGKKKERGS GKKPESAAGS QSPALPPQLK EMKSQESAAG SKLVLCRCETS SEYSSLRFKW FKNGNELNRK NKPQNIKIQQ KPGKSELRIN KASLADSGEY MCKVISKLGN DSASANITIV ESNEITGMP ASTEGAYVSS ESPIRISVST EGANTSSSTS TSTTGTSHLV KCAEKEKTFC VNGGECFMVK DLSNPSRYLC KCPNEFTGDR CQNYVMASFY KHLGIEFMEA EELYQK
Characteristics:	Recombinant Human Pro-neuregulin-1 is produced by our E. coli expression system. The target protein is expressed with sequence (Met1-Lys246) of Human NRG1.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered
Endotoxin Level:	Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test

Target Details

Target: NRG1-beta 1

Abstract: [NRG1-beta 1 Products](#)

Background: Neuregulin-1 (heregulin-1, NRG1) is a member of neuregulin family, which is comprised of four genes that encode a large number of secreted or membrane-bound isoforms. All family members share an EGF-like domain that interacts with the ErbB family of tyrosine kinase receptors. NRG1 isoforms can be classified into type I, type II and type III isoforms. NRG1 directs ligand for ERBB3 and ERBB4 tyrosine kinase receptors, concomitantly recruits ERBB1 and ERBB2 coreceptors, resulting in ligand-stimulated tyrosine phosphorylation and activation of the ERBB receptors. NRG proteins show distinct spatial and temporal expression patterns and play important roles during development of both the nervous system and the heart.

Synonyms: Pro-neuregulin-1, Neuregulin-1 beta 1, NRG1-beta 1, HRG1-beta 1 ECD, EGF, NRG1, GGF, HGL, HRGA, NDF, SMDF,

Molecular Weight: 29 kDa

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL.
Dissolve the lyophilized protein in ddH₂O.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Buffer: Lyophilized from a 0.2 µm filtered solution of 4 mM HCl.

Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

Storage: 4 °C/-20 °C/-80 °C

Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7°C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20°C for 3 months.
