

Datasheet for ABIN2116706
anti-LDHB antibody (AA 1-334)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	LDHB
Binding Specificity:	AA 1-334
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This LDHB antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Lactate Dehydrogenase B (LDHB)
Immunogen:	RPB698Hu01Recombinant Lactate Dehydrogenase B (LDHB)
Sequence:	MGHHHHHHSG SEF- MATLKEKLI PVAEEEEATVP NNKITVVGVG QVGMACAISI LGKSLADELA LVDVLEDKLG GEMMDLQHGS LFLQTPKIVA DKDYSVTANS KIVVVTAGVR QQEGESRLNL VQRNVNVFKF IIPQIVKYSP DCIIIVVSNP VDILTYVTWK LSGLPKHRVI GSGCNLDSAR FRYLMAEKLK IHPSSCHGWI LGEHGDSSVA VWSGVNVAGV SLQELNPEMG TDNDSSENWKE VHKMVVESAY EVIKLKGYN WAIGLSVADL IESMLKNLSR IHPVSTMVKG MYGIENEVFL SLPCILNARG LTSVINQKLK DDEVAQLKKS ADTLWDIQKD LKDL
Clone:	C7
Isotype:	IgG

Product Details

Specificity:	The antibody is a mouse monoclonal antibody raised against LDHB. It has been selected for its ability to recognize LDHB in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	LDHB
Alternative Name:	LDHB (LDHB Products)
Background:	LDH-B, LDH-H, Renal carcinoma antigen NY-REN-46
Pathways:	Warburg Effect

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling

Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1.