antibodies -online.com





anti-P2RY8 antibody (C-Term)

2 Images



Go to Product page

Overview

Quantity:	50 μg
Target:	P2RY8
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This P2RY8 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 18 amino acid peptide from C-terminal cytoplasmic domain of human P2RY8 / P2Y8. Percent identity with other species by BLAST analysis: Human (100%), Gorilla (94%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human P2RY8 / P2Y8. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human (100%) Gorilla (94%).
Purification:	Immunoaffinity purified

Target Details

Target:	P2RY8
Alternative Name:	P2RY8 / P2Y8 (P2RY8 Products)
Background:	Name/Gene ID: P2RY8
	Subfamily: Orphan-A
	Family: GPCR
	Synonyms: P2RY8, p2RY8, p2Y purinoceptor 8, p2Y8
Gene ID:	286530

Application Details

Ann	lication	Notes:
\neg pp	iication	INOLUS.

Approved: ELISA, IHC, IHC-P (20 µg/mL)

Usage: Immunohistochemistry: This antibody was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for this antibody was determined to be 20 μ g/mL.

Comment:

Target Species of Antibody: Human

Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for

up to 1 month. Avoid freeze-thaw cycles.

Expiry Date:


12 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Human Brain, Cortex (formalin-fixed, paraffinembedded) stained with P2RY8 antibody ABIN213464 at 20 ug/ml followed by biotinylated goat anti-rabbit IgG secondary antibody ABIN481713, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry

Image 2. Anti-P2RY8 / P2Y8 antibody IHC of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval.