antibodies - online.com







anti-GPR32 antibody (C-Term)

Images



\sim			
	IV/E	۱//۱۲	$I \cap V$

Quantity:	50 μg
Target:	GPR32
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR32 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 15 amino acid peptide from C-terminus of human GPR32. Percent identity with other species by BLAST analysis: Human (100%), Marmoset (87%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human GPR32. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FPR3 (53 $\%$).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human (100%) Marmoset (87%).
Purification:	Immunoaffinity purified

Target Details

Target:	GPR32
Alternative Name:	GPR32 (GPR32 Products)
Background:	Name/Gene ID: GPR32
	Subfamily: Orphan-A
	Family: GPCR
	Synonyms: GPR32, G protein-coupled receptor 32, Resolvin D1 receptor, RVDR1
Gene ID:	2854

Application Details

Application Notes:	Approved: ELISA, IHC, IHC-P (20 µg/mL)
, .pp.,.oato	, pp. 5 - 5 a. 22 - 5 , 5 (25 pg, 2)

Usage: Immunohistochemistry: This antibody was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for this antibody was determined to be 20 μ g/mL.

|--|--|--|

Restrictions: For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C fo

up to 1 month. Avoid freeze-thaw cycles.

Expiry Date:


12 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Human Brain, Cerebellum (formalin-fixed, paraffinembedded) stained with GPR32 antibody ABIN213488 at 20 ug/ml followed by biotinylated goat anti-rabbit IgG secondary antibody ABIN481713, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry

Image 2. Anti-GPR32 antibody IHC of human brain, cerebellum. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.