

## Datasheet for ABIN2144933

## anti-CD57 antibody (PE)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	25 tests
Target:	CD57 (B3GAT1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD57 antibody is conjugated to PE
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Functional Studies (Func)

## **Product Details**

4AHNK-1

Clone:

Isotype:	IgM
Characteristics:	4AHNK-1 reacts with HNK1 Molecule, a 110 kDa carbohydrate antigen associated with myelin-
	associated glycoprotein. CD57 expressed on 7-35 % of normal peripheral blood lymphocytes
	including a subset of naturel killer cells, a subset of CD8+ peripheral blood suppressor /
	cytotoxic T cells, and on some neural tissues. HNK is not expression on granulocytes, platelets,
	red blood cells and thymocytes

## Target Details

Target:	CD57 (B3GAT1)
Alternative Name:	CD57 (B3GAT1 Products)

Precaution of Use:

Storage:

Pathways:

Glycosaminoglycan Metabolic Process

Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Assay Procedure:	<ul> <li>Take 100 µL peripheral blood anticoagulated by EDTA and add to the bottom of 5 mLtube,</li> <li>Add 10 µL labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light,</li> <li>Add 2 ml1×RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells (recommended: RBC lysing Solution 10×,Cat.: FXP001),</li> <li>Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant,</li> <li>Add 2 mLPBS wash buffer to resuspend the cells, then1000 rpm centrifugation for 5 minutes, discard the supernatant,</li> <li>Add 0.5 mLPBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).</li> <li>[PBS wash buffer: PBS +1 % FBS +0.1 % NaN3, Cat.: FXP005]</li> <li>[Cell fixation: 2 % formaldehyde solution]</li> </ul>	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Phosphate-buffered solution, pH 7.4, containing and 0.2 % (w/v) BSA	
Preservative:	Sodium azide	

should be handled by trained staff only.

4°C

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which