

Datasheet for ABIN2145411

Mouse IgM isotype control (PE)





Overview

Overview	
Quantity:	100 tests
Target:	IgM
Host:	Mouse
Conjugate:	PE
Application:	Isotype Control (IsoC), Flow Cytometry (FACS), Immunofluorescence (IF)
Product Details	
Isotype:	IgM
Characteristics:	This antibody can be used as a mouse IgM isotype control in flow cytometry and other applications.
Target Details	
Target:	IgM
Abstract:	IgM Products
Target Type:	Antibody
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Assay Procedure:	 Take 100 µL peripheral blood anticoagulated by EDTA and add to the bottom of 5 mLtube, Add 20 µL labeled antibody to the bottom of flow tube mixing with the whole blood, incubate


- Add 2 ml1×RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells (recommended: RBC lysing Solution 10×,Cat.: FXP001),
- Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant,
- Add 2 mLPBS wash buffer to resuspend the cells, then1000 rpm centrifugation for 5 minutes, discard the supernatant,
- Add 0.5 mLPBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).
- [PBS wash buffer: PBS +1 % FBS +0.1 % NaN3, Cat.: FXP005]
- [Cell fixation: 2 % formaldehyde solution]

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Phosphate-buffered solution, pH 7.4, containing and 0.2 % (w/v) BSA
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C

Images



Flow Cytometry

Image 1.