antibodies -online.com







anti-BRF2 antibody (AA 151-250)



Image



Overview

Quantity:	100 μL
Target:	BRF2
Binding Specificity:	AA 151-250
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRF2 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details


Immunogen:	KLH conjugated synthetic peptide derived from human BRF2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Pig
Purification:	Purified by Protein A.
T	

Target Details

Target: BRF2

Target Details	
Alternative Name:	Brf2 (BRF2 Products)
Background:	Synonyms: ZFP36L2, BRF2, Butyrate response factor 2, EGF response factor 2, ERF 2, ERF2,
	Protein TIS11D, RNF162C, zinc finger protein 36, C3H type like 2, BRF2_HUMAN.
	Background: TIS11D is a member of the TIS11 family of early response genes. Family members
	are induced by various agonists such as the phorbol ester TPA and the polypeptide mitogen
	EGF. The encoded protein contains a distinguishing putative zinc finger domain with a
	repeating cys-his motif. This putative nuclear transcription factor most likely functions in
	regulating the response to growth factors.
Gene ID:	678
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



ELISA

Image 1. Antigen: 0.2ug/100ul Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG-HRP) at 1: 5000,TMB staining,Read the data in MicroplateReader by 450