# antibodies - online.com







# anti-FAT4 antibody (AA 4351-4450) (AbBy Fluor® 488)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	FAT4
Binding Specificity:	AA 4351-4450
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAT4 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CDHF14
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	FAT4
Alternative Name:	Cdhf14 (FAT4 Products)

#### **Target Details**

#### Background:

Synonyms: Cadherin family member 14, CDHF14, FAT tumor suppressor homolog 4, Fat-like cadherin protein FAT-J, FAT4, FAT4\_HUMAN, FATJ, hFat4, Nbla00548, Protocadherin Fat 4. Background: The cadherins represent a family of Ca2+-dependent adhesion molecules that function to mediate cell to cell binding that is critical for the maintenance of structure and morphogenesis. Cadherins contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short C-terminal intracellular domain interacts with a variety of cytoplasmic proteins, including beta-catenin, to regulate cadherin function. The cadherin superfamily includes cadherins, protocadherins, desmogleins and desmocollins. FAT4 (FAT tumor suppressor homolog 4), also known as FAT-J, CDHF14 or CDHR11, is a 4,981 amino acid single-pass type I membrane protein that belongs to the protocadherin subfamily of cadherins and localizes to the primary cilia of kidney. Widely expressed, FAT4 contains thirty-four cadherin domains, six EGF-like domains and two laminin G-like domains. FAT4 may function in the regulation of planar cell polarity.

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months