# antibodies .- online.com







# anti-CEP120 antibody (AA 351-450) (HRP)



| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | CEP120  |  |
| Binding Specificity: | AA 351-450  |  |
| Reactivity:          | Human, Mouse  |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This CEP120 antibody is conjugated to HRP   |  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human CEP120 |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Cross-Reactivity:     | Human, Mouse   |  |
| Predicted Reactivity: | Rat  |  |
| Purification:         | Purified by Protein A.                                     |  |

## Target Details

| Target:           | CEP120                   |
|-------------------|--------------------------|
| Alternative Name: | CEP120 (CEP120 Products) |

### **Target Details**

| 0                   |   |  |
|---------------------|---|--|
| Background:         | Synonyms: CCDC100, Centrosomal protein of 120 kDa, Cep120, Coiled-coil domain-containing        |  |
|                     | protein 100   |  |
|                     | Background: Plays a role in the microtubule-dependent coupling of the nucleus and the           |  |
|                     | centrosome. Involved in the processes that regulate centrosome-mediated interkinetic nuclear    |  |
|                     | migration (INM) of neural progenitors and for proper positioning of neurons during brain        |  |
|                     | development. Also implicated in the migration and selfrenewal of neural progenitors. May play a |  |
|                     | role in centriole duplication during mitosis (By similarity).                                   |  |
| Gene ID:            | 153241  |  |
| UniProt:            | Q8N960  |  |
| Application Details |   |  |
| Application Notes:  | WB 1:300-5000   |  |
|                     | IHC-P 1:200-400   |  |
|                     | IHC-F 1:100-500   |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 μg/μL   |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |  |
|                     | 50 % Glycerol.  |  |
| Preservative:       | ProClin   |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |  |
|                     | handled by trained staff only.  |  |
| Handling Advice:    | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish        |  |
|                     | peroxidase.   |  |
| Storage:            | -20 °C  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.               |  |
| Expiry Date:        | 12 months   |  |
|                     |   |  |