antibodies .- online.com







anti-GADD34 antibody (AA 401-500) (Alexa Fluor 488)



| \sim | |
|--------|----------|
| ()\/△ | rview |
| \cup | 1 410 44 |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | GADD34 (PPP1R15A) |
| Binding Specificity: | AA 401-500 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GADD34 antibody is conjugated to Alexa Fluor 488 |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GADD34 |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GADD34 (PPP1R15A) |
|-------------------|---|
| Alternative Name: | GADD34 (PPP1R15A Products) |
| Background: | Synonyms: GADD34, Protein phosphatase 1 regulatory subunit 15A, Growth arrest and DNA |

Storage Comment:

12 months

Expiry Date:

| Target Details | |
|---------------------|--|
| | damage-inducible protein GADD34, Myeloid differentiation primary response protein MyD116 |
| | homolog, PPP1R15A |
| | Background: Recruits the serine/threonine-protein phosphatase PP1 to dephosphorylate the |
| | translation initiation factor eIF-2A/EIF2S1, thereby reversing the shut-off of protein synthesis |
| | initiated by stress-inducible kinases and facilitating recovery of cells from stress. Down- |
| | regulates the TGF-beta signaling pathway by promoting dephosphorylation of TGFB1 by PP1. |
| | May promote apoptosis by inducing TP53 phosphorylation on 'Ser-15'. |
| Gene ID: | 23645 |
| UniProt: | 075807 |
| Pathways: | ER-Nucleus Signaling, Unfolded Protein Response |
| Application Details | |
| Application Notes: | FCM 1:20-100 |
| | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.