antibodies -online.com





Rabbit anti-Dog IgG Antibody (Cy7)



| _ | | | | | | | |
|---|---|---|----|----|---|---|---|
| 0 | V | е | r١ | /1 | е | V | 1 |

| Quantity: | 200 μL | |
|-----------------|---|--|
| Target: | IgG | |
| Reactivity: | Dog | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | Cy7 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| Isotype: | IgG | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | IgG | |
| Abstract: | IgG Products | |
| Target Type: | Antibody | |
| Background: | Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms | |

The IgG molecule has two separate functions, to bind to the pathogen that elicited the response

Target Details

and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.

Application Details

| Application Notes: | IF(IHC-P): (1:500-2000), IF(IHC-F): (1:500-2000), IF(ICC): (1:500-1000) Optimal working dilution should be determined by the investigator. |
|--------------------|---|
| Comment: | Exitation/Emission: 743nm/767nm |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 100 μg/mL BSA, 50 % glycerol and 0.09 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at 4 °C for 12 months. |