# antibodies -online.com





# **Goat anti-Cow IgG Antibody (Cy3)**



( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Quantity:	200 μL
Target:	IgG
Reactivity:	Cow
Host:	Goat
Clonality:	Polyclonal
Conjugate:	СуЗ
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Isotype:	IgG
Purification:	Purified by Protein A.

#### **Target Details**

l arget:	IgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels
	between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com

and the molecules are produced by B lymphocytes as a part of our adaptive immune response.

The IgG molecule has two separate functions, to bind to the pathogen that elicited the response

#### **Target Details**

and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.

# **Application Details**

Application Notes:	IF(IHC-P): (1:500-2000), IF(IHC-F): (1:500-2000), IF(ICC): (1:500-1000)
	Optimal working dilution should be determined by the investigator.
Comment:	Exitation/Emission: 512,550nm/570,615nm
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 100 μg/mL BSA, 50 % glycerol and 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at 4 °C for 12 months.