



[Go to Product page](#)

Datasheet for ABIN2176670

Rabbit anti-Monkey IgG Antibody (Cy7)

Overview

| | |
|--------------|---|
| Quantity: | 200 µL |
| Target: | IgG |
| Reactivity: | Monkey |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | Cy7 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|---------------|------------------------|
| Isotype: | IgG |
| Purification: | Purified by Protein A. |

Target Details

| | |
|--------------|--|
| Target: | IgG |
| Abstract: | IgG Products |
| Target Type: | Antibody |
| Background: | Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions, to bind to the pathogen that elicited the response |

Target Details

and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.

Application Details

Application Notes: IF(IHC-P): (1:500-2000), IF(IHC-F): (1:500-2000), IF(ICC): (1:500-1000)
Optimal working dilution should be determined by the investigator.

Comment: Excitation/Emission: 743nm/767nm

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 100 µg/mL BSA, 50 % glycerol and 0.09 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at 4 °C for 12 months.