# antibodies - online.com







## anti-IL1RAP antibody (AA 251-350)



Image



#### Overview

Quantity:	100 μL
Target:	IL1RAP
Binding Specificity:	AA 251-350
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL1RAP antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human IL1RAP/IL1R3
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat
Purification:	Purified by Protein A.

### Target Details

Target:	IL1RAP
Alternative Name:	IL1RAP (IL1RAP Products)
Background:	Synonyms: IL1R3, C3orf13, IL-1RAcP, Interleukin-1 receptor accessory protein, IL-1 receptor


	accessory protein, Interleukin-1 receptor 3, IL-1R-3, IL-1R3, IL1RAP
	Background: Coreceptor for IL1RL2 in the IL-36 signaling system (By similarity). Coreceptor
	with IL1R1 in the IL-1 signaling system. Associates with IL1R1 bound to IL1B to form the high
	affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-
	kappa-B and other pathways. Signaling involves the recruitment of adapter molecules such as
	TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor
	subunits. Recruits TOLLIP to the signaling complex. Does not bind to interleukin-1 alone,
	binding of IL1RN to IL1R1, prevents its association with IL1R1 to form a signaling complex. The
	cellular response is modulated through a non-signaling association with the membrane IL1R2
	decoy receptor. Secreted forms (isoforms 2 and 3) associate with secreted ligand-bound IL1R2
	and increase the affinity of secreted IL1R2 for IL1B, this complex formation may be the
	dominant mechanism for neutralization of IL1B by secreted/soluble receptors.
Gene ID:	3556
UniProt:	Q9NPH3
Pathways:	NF-kappaB Signaling, Growth Factor Binding
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400

Handli	ng
--------	----

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

For Research Use only



#### **Western Blotting**

**Image 1.** Mouse liver lysates probed with Rabbit Anti-IL1RAP Polyclonal Antibody, Unconjugated (ABIN2176996) at 1:300 overnight at 4 °C. Followed by conjugation to secondary antibody at 1:5000 for 90 min at 37 °C.